

AD A049986

AD No. 1  
DDC FILE COPY

✓ARO 14435.7-M

(5)

DISTRIBUTION OF A SUM OF WEIGHTED CHI-SQUARE VARIABLES

BY

H. SOLOMON and M. A. STEPHENS

See back  
page for 1473

TECHNICAL REPORT NO. 5

JUNE 2, 1977

PREPARED UNDER GRANT

DAAG29-77-G-0031

FOR THE U.S. ARMY RESEARCH OFFICE

DDC  
PREPARED  
FEB 15 1978  
F

Reproduction in Whole or in Part is Permitted  
for any purpose of the United States Government

Approved for public release; distribution unlimited.

DEPARTMENT OF STATISTICS

STANFORD UNIVERSITY

STANFORD, CALIFORNIA



(18) ARO (19) 14435.7M

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER (14) TR-245, TR-5	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) (6) Distribution of a Sum of Weighted Chi-Square Variables.	5. TYPE OF REPORT & PERIOD COVERED (9) Technical Report	
7. AUTHOR(s) (10) H./Solomon M. A./Stephens	6. PERFORMING ORG. REPORT NUMBER	
8. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics Stanford University Stanford, California 94305	9. CONTRACT OR GRANT NUMBER(s) (15) N00014-76-C-0475 DAAG29-77-G-0031	
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Research Office Post Office Box 12211 Research Triangle Park, NC 27709	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS P-14435-M	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE (11) June 2, 1977 13. NUMBER OF PAGES 12 15. SECURITY CLASS. (of this report) (12) 16p. 16. DECLASSIFICATION/CONTROLLING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents. This report partially supported under Office of Naval Research Contract N00014-76-C-0475 (NR-042-267) and issued as Technical Report No. 245		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Quadratic forms; weighted chi-square variables; Imhof technique		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  (see reverse side)		

DD FORM 1473  
1 JAN 73

EDITION OF 1 NOV 69 IS OBSOLETE  
S/N 0102-010-6001

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

332 580

Yun

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20. In this paper we discuss how to obtain percentage points of the distribution of a sum of  $k$  weighted chi-square variables. A table of points is given for selected weights, for  $k = 6, 8, 10$  (existing tables are for  $k \leq 5$ ). The new points have been calculated by a technique of Imhof (1961) which gives very accurate values but is expensive in computer time. Therefore two new approximations are considered: a Pearson curve fit using four moments and a  $\chi^2$  fit using three. Both give excellent results in the upper tail, the tail usually needed for practical applications; the  $\chi^2$  fit appears superior to other approximations in the lower tail.

chi-square

245/5

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

DISTRIBUTION OF A SUM OF WEIGHTED CHI-SQUARE VARIABLES

by

H. Solomon and M. A. Stephens


TECHNICAL REPORT NO. 5

JUNE 2, 1977

Prepared under Grant DAAG29-77-G-0031

For the U. S. Army Research Office

Herbert Solomon, Project Director



Approved for public release; distribution unlimited.

DEPARTMENT OF STATISTICS  
STANFORD UNIVERSITY  
STANFORD, CALIFORNIA

Partially supported under Office of Naval Research Contract N00014-76-C-0475  
and issued as Technical Report No. 245.

THE FINDINGS IN THIS REPORT ARE NOT TO BE  
CONSTRUED AS AN OFFICIAL DEPARTMENT OF  
THE ARMY POSITION, UNLESS SO DESIGNATED  
BY OTHER AUTHORIZED DOCUMENTS.

## Distribution of a Sum of Weighted Chi-Square Variables

### 1. INTRODUCTION

Distributions of quadratic forms of the type

$$(1) \quad Q_k = \sum_{j=1}^k c_j (x_j + a_j)^2$$

where  $x_j$  are i.i.d. standard normal variables (i.e., mean 0 and variance 1), and where  $c_j$  and  $a_j$  are non-negative constants, arise in many problems in statistics. A brief survey of such problems, with references, is in Jensen and Solomon (1972). We employ their notation in this report. An important class of problems, where there has been much recent activity, concerns asymptotic theory of goodness of fit tests. Statistics related to Pearson's chi-square, when unknown parameters are estimated by various methods, and when cells are allowed to be data-dependent, have asymptotic distributions of the  $Q_k$  type; for a survey see Moore (1976). Statistics based on the empirical distribution function (EDF statistics) also have asymptotic distributions of this type; references are given below.

Exact significance points of  $Q_k$ , for selected  $c_k$ , and all  $a_k = 0$ , have been published, for  $k = 2$  and 3, by Grad and Solomon (1955) and by Solomon (1960), reproduced in Owen (1962), and for  $k = 4$  and 5, by Johnson and Kotz (1968); Solomon gives, also, tables of  $P_k(t) = P(Q_k < t)$ , for certain values of  $t$ . The mathematical difficulties involved in obtaining exact values increase with  $k$ , and several approximate methods have been suggested. The most accurate, as we show later, appears to be that of Imhof (1961), in which the characteristic function of the  $Q_k$  distribution is inverted numerically to give  $P_k(t)$  for given  $t$ . In the calculations, an integral with an infinite upper limit must be calculated for each  $t$ . Imhof gives a bound on the accuracy obtained when this upper

ACCESSION FOR	NTIS	DDC	UNIVERSITY MICROFILMS	BY	DISTRIBUTION STATEMENTS	DATE

1

limit is replaced by a finite  $T$ . This is apart from inaccuracies in the method of numerical integration. Thus a measure of accuracy can be attained. The Imhof method has been used by Durbin and Knott (1972), Durbin, Knott and Taylor (1975) and Pettitt and Stephens (1975) in approximating distributions of EDF statistics. These require  $k$  to be infinite and make use of a modification of Imhof's technique suggested by Durbin and Knott.

## 2. NEW SIGNIFICANCE POINTS FOR $Q_k$

It is a convenience to have exact points at hand whenever possible, so for selected  $c_k$ , and all  $a_k = 0$ , we give significance points of  $Q_k$ , for  $k = 6, 8, 10$ , in Table 1. These have been calculated by Imhof's method. They will also provide useful anchor points for comparison with other approximations already in the literature or to be suggested in Section 3. In Table 2, among several comparisons, we give exact values of significance points, and values calculated by Imhof's method to a high order of accuracy. It can be seen that the Imhof technique gives excellent results.

## 3. NEW APPROXIMATIONS FOR $Q_k$

There is clearly a need for an approximation to the  $Q_k$  distribution for problems where  $a_k$  do not appear in existing Tables; such would be the case for example, in the distribution theory of statistics of the chi-square type referred to above. The Imhof method can be almost regarded as exact, but it does not adapt easily to step-by-step increases in  $t$

such as are needed for good inverse interpolation to obtain significance points, and when high accuracy is sought, the authors have found it relatively expensive in computer time. The need for another good approximation therefore exists. Several authors have searched for such approximations and comparisons and references are given in Jensen and Solomon (1972).

Jensen and Solomon also give an approximation of the Wilson-Hilferty type which appears to be one of the most effective. Their approximation takes  $z = (Q_k/\theta_1)^h$ , where  $\theta_1$  is the mean of  $Q_k$ , and approximates  $z$  by a normal distribution where the mean and variance of the approximating distribution depend on the first three cumulants of  $Q_k$ .

We now propose two additional approximations: (1) fitting a Pearson curve with the same first four moments as  $Q_k$ ; and (2) fitting  $Q_k = Aw^r$ , where  $w$  has the  $\chi_p^2$  distribution, and where  $A, r, p$  are determined by the first three moments of  $Q_k$ .

Previous authors (Patnaik (1949), Pearson (1959)) have fitted Pearson curves using less than four moments. Considerable experience with four-moment Pearson curve fitting suggests this method will be very accurate in the long upper tail of  $Q_k$  (the tail required for most applications). A practical drawback is that, once the moments have been calculated, interpolation is necessary, either in the tables of Johnson, Nixon, Amos and Pearson (1963), also reproduced in Biometrika Tables for Statisticians, Volume 2 (Pearson and Hartley (1972)), or in extensions of these tables recently produced by Amos and Daniel (1971) and by Bouwer and Bargmann (1973).

The Pearson curve fits will be much less effective in the lower tail;



matching the moments does not even insure that the value of  $P_k(0) = 0$ , i.e., that the distribution of  $Q_k$  "starts" at zero. This weakness is shared by other approximations, including that of Jensen and Solomon. However, approximation (2) above does automatically start at zero. These considerations suggest it will be better as an approximation, at least in the lower tail.

### 3.1 The three-moment chi-square fit

The distribution of  $Q_k$  is to be fitted by  $Q_k = Aw^r$ , where  $w$  has the  $\chi_p^2$  distribution and the constants  $A, p, r$  will be found by matching moments. Let  $p/2 = v$ , and let  $C = \Gamma(v)$ ; the moments of  $z$  are then

$$\mu = E(z) = A 2^r \{\Gamma(r+v)\}/C,$$

$$\mu_2' = A^2 4^r \{\Gamma(2r+v)\}/C,$$

$$\mu_3' = A^3 8^r \{\Gamma(3r+v)\}/C.$$

Define

$$R_2 = \mu_2'/\mu^2 = C \Gamma(2r+v)/\{\Gamma(r+v)\}^2$$

and

$$R_3 = \mu_3'/\mu^3 = C^2 \Gamma(3r+v)/\{\Gamma(r+v)\}^2.$$

Given  $R_2$  and  $R_3$ , these equations can be solved for  $r$  and  $v$  and then  $A$  is obtained from the expression for  $\mu$ . Computer routines are available to perform these operations and then to calculate probabilities or significance points of  $\chi^2$ , even with non-integer degrees of freedom. Significance points for  $\chi^2$  with degrees of freedom differing by 0.2

are given in Pearson and Hartley (1972).

The  $\chi^2$  approximation is not sensitive to employing for  $p$  the closest integer to  $p$ . If  $R_2$  is then used to solve for  $r$ , a good approximation will often result. Further details on this method of approximation, with other areas of application, will be given in a later report.

### 3.2 Accuracy of various approximations

Table 2 compares significance points for several distributions  $Q_k$  for which exact (E) values exist in Solomon (1960) or Johnson and Kotz (1968); the approximations are Imhof's (I), the Jensen-Solomon approximation (J), and the new approximations, the four-moment Pearson curve fit (P) and the new three-moment  $\chi^2$  fit (S). In Table 3 these comparisons are continued for higher values of  $k$ , for which no exact values exist. In these tables, the values of  $c_k$  have been chosen to sum either to 1 or to  $k$ , to enable comparisons to be made directly with values given in the references. Jensen and Solomon compare values of  $P_k(k)$  rather than significance points, so in Table 4 we give some comparisons of this type. In Tables 2, 3, and 4, values of  $\sqrt{\beta_1}$  and  $\beta_2$  are given in square brackets beneath the values of  $c_k$ , where  $\beta_1 = \mu_3/\mu_2^3$  is a measure of skewness and  $\beta_2 = \mu_4/\mu_2^2$  is a measure of kurtosis.

### 3.3 Comments

In comparing approximations with exact values, the important quantity is  $\alpha'$ , the exact significance level realized by an approximate point calculated for significance level  $\alpha$ . The values in Table 2 show immediately the excellent accuracy of the Imhof technique, and values in Table 1 obtained by Imhof's method, and quoted again in Table 3, may for practical

purposes be regarded as accurate. Apart from this, all three approximations (P, J, S) generally give excellent accuracy in the upper tail. In the lower tail, the Pearson curves often are very poor, especially for higher values of  $\sqrt{\beta_1}$  and  $\beta_2$ ; Tables 2 and 3 show the clear supremacy of S over P. Table 4 suggests that S is more accurate than J in both tails; however, all approximations (except Imhof's) become relatively less good in the lower tail as the skewness and kurtosis of  $Q_k$  increase. Jensen and Solomon have already demonstrated an overall supremacy of J over other approximations discussed by them. From the picture presented here S is as good as or better than J and this merits consideration as an approximation if only three moments are to be used and especially for accuracy in the lower tail. Probability levels for S are easily obtained all along the curve and this is useful in combining significance tests; one of the occasions when the lower tail becomes important.

### References

- Amos, D. E. and Daniel, S. L. (1971). Tables of percentage points of standardized Pearson distributions. Research Report SC-RR-71 0348, Albuquerque, New Mexico: Sandia Laboratories.
- Bouwer, H. and Bargmann, R. E. (1973). Tables of the standardized percentage points of the Pearson system of curves in terms of  $\beta_1$  and  $\beta_2$ . Technical Report No. 107, Department of Statistics and Computer Science, University of Georgia.
- Durbin, J. and Knott, M. (1972). Components of Cramér-von Mises statistics, I. J. R. Statist. Soc. B, 34, 290-307.
- Durbin, J., Knott, M. and Taylor, C. C. (1975). Components of Cramér-von Mises statistics, II. J. R. Statist. Soc. B, 37, 216-237.
- Grad, A. and Solomon, H. (1955). Distribution of quadratic forms and some applications. Annals of Mathematical Statistics, 26, 464-477.
- Imhof, J. P. (1961). Computing the distribution of quadratic forms in normal variables. Biometrika, 48, 417-426.
- Jensen, D. R. and Solomon, H. (1972). A Gaussian approximation to the distribution of a definite quadratic form. J. Amer. Statist. Ass., 67, 898-902.
- Johnson, N. L. and Kotz, S. (1968). Tables of distributions of positive definite quadratic forms in central normal variables. Sankhyā B, 30, 303-314.
- Johnson, N. L., Nixon, E., Amos, D. E. and Pearson, E. S. (1963). Table of percentage points of Pearson curves, for given  $\sqrt{\beta_1}$  and  $\beta_2$ , expressed in standard measure. Biometrika, 50, 459-498.
- Moore, D. S. (1976). Recent developments in Chi-square tests for goodness-of-fit. Technical Report 459, Department of Statistics, Purdue University.
- Owen, D. B. (1962). Handbook of Statistical Tables. Reading, Massachusetts: Addison-Wesley.
- Patnaik, P. B. (1949). The noncentral  $\chi^2$  and F-distributions and their applications. Biometrika, 36, 202-232.
- Pearson, E. S. (1959). Note on an approximation to the distribution of non-central  $\chi^2$ . Biometrika, 46, 364.
- Pearson, E. S. and Hartley, H. O. (1972). Biometrika Tables for Statisticians, Vol. 2. Cambridge: Cambridge University Press.
- Pettitt, A. N. and Stephens, M. A. (1976). Modified Cramer-von Mises statistics for censored data. Biometrika, 63, 291-298.

Solomon, H. (1960). Distribution of quadratic forms--tables and applications. Applied Mathematics and Statistics Laboratories, Technical Report 45, Stanford University.

Solomon, H. and Stephens, M. A. (1975). Pearson curves revisited. Technical Report No. 226, Department of Statistics, Stanford University, Stanford, California.

TABLE 1

Percentage Points for  $\chi^2_k$  — All  $a_j$  are Zero

	c-values	$\alpha$									
		.01	.025	.05	.10	.25	.75	.90	.95	.975	.99
k = 6	.2 .2 .2 .2 .1 .1	.1354	.1974	.2634	.3559	.5613	1.3077	1.8001	2.1473	2.4834	2.9135
	.3 .2 .2 .1 .1 .1	.1335	.19	.2522	.3424	.5448	1.3046	1.8309	2.2140	2.5922	3.0699
	.3 .3 .2 .1 .05 .05	.1162	.1670	.2242	.3090	.5073	1.3157	1.9023	2.3326	2.7573	3.3140
	.3 .3 .1 .1 .1 .1	.1240	.1814	.2429	.3299	.5277	1.3025	1.8651	2.2828	2.6984	3.2513
	.4 .3 .15 .05 .05 .05	.1044	.1506	.2032	.2821	.4722	1.3145	1.9695	2.4641	2.9610	3.6224
	.4 .25 .15 .1 .05 .05	.1127	.1619	.2174	.2997	.4931	1.3064	1.9277	2.3998	2.8776	3.5195
	.4 .2 .2 .1 .05 .05	.1139	.1636	.2196	.3027	.4973	1.3065	1.9178	2.3816	2.8523	3.4875
	.4 .2 .1 .1 .1 .1	.1215	.1778	.2381	.3235	.5178	1.2923	1.8787	2.3315	2.7988	3.4299
	.5 .3 .05 .05 .05 .05	.0906	.1310	.1774	.2481	.4243	1.3126	2.0655	2.6489	3.2436	4.0451
	.5 .2 .1 .1 .05 .05	.1053	.1512	.2032	.2804	.4637	1.2983	1.9851	2.5440	3.1264	3.9231
	.5 .1 .1 .1 .1 .1	.1126	.1649	.2211	.3010	.4850	1.2768	1.9455	2.5000	3.0857	3.6832
	.6 .1 .1 .1 .05 .05	.0937	.1383	.1868	.2570	.4265	1.2625	2.0729	2.7493	3.4534	4.4221
	.2 .2 .2 .2 .05 .05 .05	.1717	.2306	.2930	.3814	.5763	1.2933	1.2933	2.12	2.4521	2.8799
	.2 .2 .2 .1 .1 .05 .05	.1842	.2460	.3108	.4014	.5966	1.2844	1.7386	2.0614	2.3745	2.7794
	.2 .2 .1 .1 .1 .1 .1	.1970	.2618	.3288	.4212	.6168	1.2771	1.7007	1.996	2.2895	2.6649
	.3 .3 .1 .1 .05 .05 .05	.1597	.2147	.2731	.3563	.5428	1.2862	1.8424	2.2592	2.6753	3.2252
k = 8	.3 .2 .2 .1 .05 .05 .05	.1656	.2225	.2829	.3688	.5596	1.2891	1.8075	2.1878	2.5646	3.1613
	.3 .2 .1 .1 .1 .05 .05	.1779	.2378	.3006	.3887	.5801	1.2785	1.7674	2.1298	2.4930	2.9777
	.4 .3 .05 .05 .05 .05 .05	.1404	.1894	.2420	.3182	.4950	1.2912	1.9381	2.4313	2.9279	3.5896
	.4 .2 .1 .1 .05 .05 .05	.1572	.2111	.2685	.3502	.5333	1.2751	1.8563	2.3096	2.7764	3.4116
	.5 .2 .05 .05 .05 .05 .05	.1368	.1843	.2353	.3090	.4795	1.2700	1.9659	2.5263	3.1099	3.9082
	.5 .1 .1 .1 .05 .05 .05	.1477	.1983	.2521	.3286	.5012	1.2510	1.9239	2.4827	3.0693	3.8717
	.2 .2 .2 .1 .05 .05 .05 .05	.2138	.2763	.3409	.4284	.6125	1.2691	1.7122	2.0291	2.3387	2.7415
	.2 .2 .1 .1 .1 .05 .05 .05	.2266	.2915	.3583	.4480	.6331	1.2619	1.6716	1.9649	2.2506	2.6192
	.2 .1 .1 .1 .1 .1 .05 .05	.2399	.3070	.3758	.4675	.6535	1.2571	1.6340	1.8975	2.1518	2.4857
	.3 .25 .1 .05 .05 .05 .05 .05	.2009	.2588	.3183	.4007	.5794	1.2656	1.7768	2.1611	2.5461	3.0563
	.3 .2 .1 .1 .05 .05 .05 .05	.2075	.2682	.3309	.4161	.5963	1.2620	1.7412	2.1013	2.4630	2.9503
	.3 .15 .1 .1 .05 .05 .05 .05	.2172	.2763	.3417	.4276	.6096	1.2571	1.7132	2.0565	2.4054	2.8773
	.4 .2 .05 .05 .05 .05 .05 .05	.1889	.2434	.2998	.3780	.5500	1.2565	1.6338	2.2866	2.7543	3.3939
	.4 .1 .1 .1 .05 .05 .05 .05	.1987	.2566	.3166	.3980	.5710	1.2429	1.7860	2.2279	2.6296	3.3357
	.5 .1 .05 .05 .05 .05 .05 .05	.1773	.2294	.2836	.3578	.5187	1.2290	1.9027	2.4648	3.0512	3.8526

TABLE 2

Comparison of Exact (E) Percentage Points and Approximations I, P, S, J (see Section 3)

c-values [ $\sqrt{\beta_1, \beta_2}$ ]		$\alpha$							
		0.01	0.025	0.05	0.10	0.90	0.95	0.975	0.99
0.5, 0.4, 0.1 [1.974, 9.000]	E	---	---	.098	.164	2.188	2.820	---	---
	I	.032	.060	.097	.164	2.187	2.818	3.450	4.290
	P	.061	.079	.109	.166	2.191	2.818	3.447	4.284
	S	.027	.053	.091	.158	2.184	2.809	3.441	4.275
0.5, 0.3, 0.2 [1.932, 9.000]	E	---	---	.110	.183	2.122	2.708	---	---
	I	.036	.068	.110	.183	2.122	2.708	3.302	4.102
	P	.056	.082	.119	.187	2.127	2.712	3.302	4.093
	S	.040	.071	.114	.186	2.125	2.713	3.308	4.103
1.5, 1.5, 0.5, 0.5 [1.771, 7.920]	J	.029	.060	.103	.175	2.132	2.740	3.365	4.217
	E	.261	.425	.627	.947	8.120	10.203	12.283	15.033
	I	.259	.424	.627	.947	8.120	10.202	12.282	15.032
	P	.387	.510	.673	.953	8.141	10.205	12.261	14.984
2.5, 0.5, 0.5, 0.5 [2.444, 12.612]	S	.252	.415	.614	.933	8.115	10.177	12.247	14.990
	J	.212	.373	.574	.894	8.152	10.287	12.456	15.385
	E	.226	.369	.545	.826	8.540	11.343	14.281	18.297
	I	.225	.368	.545	.826	8.540	11.342	14.279	18.294
1.8, 1.8, 0.6, 0.4, 0.4 [1.773, 7.957]	P	.723	---	.807	.939	8.661	11.403	---	18.133
	S	.202	.331	.495	.765	8.564	11.232	14.051	17.980
	J	.180	.295	.440	.683	8.707	11.766	15.141	20.094
	E	.448	.677	.943	1.347	9.921	12.419	14.914	18.214
3.0, 0.5, 0.5, 0.5, 0.5 [2.460, 12.750]	I	.449	.677	.943	1.347	9.921	12.419	14.913	18.213
	P	.667	.813	1.011	1.351	9.951	12.419	14.879	18.143
	S	.415	.640	.904	1.372	9.902	12.358	14.837	18.130
	J	.370	.589	.850	1.253	9.983	12.567	15.208	18.801
	E	.403	.609	.848	1.211	10.407	13.778	17.307	22.259
	I	.404	.609	.848	1.211	10.407	13.778	17.308	22.127
	P	1.104	---	1.201	1.355	10.562	13.847	---	21.919
	S	.344	.528	.750	1.103	10.428	13.591	16.958	21.735
	J	.324	.485	.680	.992	10.645	14.285	18.568	24.813

TABLE 3  
Comparison of Approximate Percentage Points

c-values [ $\sqrt{\beta_1}, \beta_2$ ]		$\alpha$							
		.01	.025	.05	.10	.90	.95	.975	.99
0.3, 0.2, 0.1, 0.1, 0.05 (6 times) [1.593, 7.380]	I	.208	.268	.331	.416	1.741	2.101	2.463	2.950
	P	.244	.289	.340	.415	1.747	2.102	2.457	2.932
	S	.213	.270	.330	.413	1.741	2.096	2.454	2.937
	J	.206	.261	.318	.398	1.771	2.159	2.558	3.107
0.4, 0.1, 0.1, 0.1, 0.05 (6 times) [2.065, 10.406]	I	.199	.257	.317	.398	1.786	2.228	2.630	3.336
	P	----	----	----	----	----	----	----	----
	S	.204	.256	.312	.389	1.794	2.214	2.655	3.274
	J	.200	.247	.296	.367	1.849	2.363	2.936	3.794
0.2, 0.2, 0.2, 0.2, 0.1, 0.1 [1.259, 5.444]	I	.135	.197	.263	.356	1.800	2.147	2.483	2.914
	P	.144	.201	.263	.355	1.801	2.148	2.482	2.912
	S	.139	.198	.263	.355	1.801	2.148	2.484	2.912
	J	.131	.192	.258	.353	1.801	2.151	2.491	2.932
0.4, 0.2, 0.1, 0.1, 0.1, 0.1 [1.828, 8.750]	I	.122	.178	.238	.324	1.879	2.332	2.799	3.429
	P	.172	.209	.255	.327	1.892	2.340	2.794	3.410
	S	.135	.186	.242	.324	1.885	2.335	2.794	3.420
	J	.128	.176	.229	.308	1.917	2.412	2.928	3.648



TABLE 4  
Comparison of Values of  $P_k(t) = P(Q_k < t)$

c-values [ $\sqrt{\beta_1, \beta_2}$ ]		t						
		0.1	0.2	0.8	1.0	2.2	3.0	5.8
0.4, 0.3, 0.3 [1.683, 7.339]	E	.0405	.1047	.5086	.6102	.9134	.9698	.9896
	J	.0449	.1065	.5077	.6105	.9143	.9697	.9893
	S	.0397	.1039	.5091	.6107	.9134	.9697	.9896
0.5, 0.4, 0.1 [1.974, 9.000]	E	.0517	.1282	.5363	.6282	.9013	.9590	.9828
	J	.0644	.1395	.5371	.6289	.9019	.9586	.9820
	S	.0566	.1328	.5335	.6257	.9019	.9595	.9831
c-values [ $\sqrt{\beta_1, \beta_2}$ ]		t						
		0.5	1.0	3.0	3.6	8.5	11.0	14.0
2.5, 0.7, 0.4, 0.4 [2.427, 12.475]	E	.0448	.1385	.5197	.6005	.8980	.9452	.9731
	J	.0645	.1742	.5415	.6153	.8944	.9407	.9686
	S	.0534	.1514	.5153	.5935	.8975	.9465	.9745
2.0, 1.0, 0.7, 0.3 [2.011, 9.647]	E	.0375	.1189	.4862	.5728	.9078	.9568	.9821
	J	.0439	.1320	.5001	.5833	.9046	.9531	.9792
	S	.0373	.1198	.4864	.5721	.9075	.9569	.9823